

## Anti-PRX II antibody



|                    |                              |
|--------------------|------------------------------|
| <b>Description</b> | Rabbit polyclonal to PRX II. |
|--------------------|------------------------------|

|                         |  |
|-------------------------|--|
| <b>Model</b>            | STJ95237   |
| <b>Host</b>             | Rabbit   |
| <b>Reactivity</b>       | Human, Mouse, Rat  |
| <b>Applications</b>     | ELISA, WB  |
| <b>Immunogen</b>        | Synthesized peptide derived from human PRX II.   |
| <b>Immunogen Region</b> | C-terminal   |
| <b>Gene ID</b>          | <a href="#">7001</a>   |
| <b>Gene Symbol</b>      | <a href="#">PRDX2</a>  |
| <b>Dilution range</b>   | WB 1:500-1:2000ELISA 1:10000   |
| <b>Specificity</b>      | PRX II Polyclonal Antibody detects endogenous levels of PRX II protein.  |
| <b>Purification</b>     | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| <b>Note</b>             | For Research Use Only (RUO).   |
| <b>Protein Name</b>     | Peroxiredoxin-2 Natural killer cell-enhancing factor B NKEF-B PRP Thiol-specific antioxidant protein TSA Thioredoxin peroxidase 1 Thioredoxin-dependent peroxide reductase 1 |
| <b>Molecular Weight</b> | 21 kDa   |
| <b>Clonality</b>        | Polyclonal   |
| <b>Conjugation</b>      | Unconjugated   |

|                              |  |
|------------------------------|--|
| <b>Isotype</b>               | IgG  |
| <b>Formulation</b>           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| <b>Concentration</b>         | 1 mg/ml  |
| <b>Storage Instruction</b>   | Store at -20°C, and avoid repeat freeze-thaw cycles.   |
| <b>Database Links</b>        | <a href="#">HGNC:9353</a> <a href="#">OMIM:600538</a>  |
| <b>Alternative Names</b>     | Peroxiredoxin-2 Natural killer cell-enhancing factor B NKEF-B PRP Thiol-specific antioxidant protein TSA Thioredoxin peroxidase 1 Thioredoxin-dependent peroxide reductase 1   |
| <b>Function</b>              | Involved in redox regulation of the cell. Reduces peroxides with reducing equivalents provided through the thioredoxin system. It is not able to receive electrons from glutaredoxin. May play an important role in eliminating peroxides generated during metabolism. Might participate in the signaling cascades of growth factors and tumor necrosis factor-alpha by regulating the intracellular concentrations of H <sub>2</sub> O <sub>2</sub> . |
| <b>Cellular Localization</b> | Cytoplasm.   |

---

**St John's Laboratory Ltd**

**F** +44 (0)207 681 2580  
**T** +44 (0)208 223 3081

**W** <http://www.stjohnslabs.com/>  
**E** [info@stjohnslabs.com](mailto:info@stjohnslabs.com)