


Anti-Rab 35 antibody



Description	Rabbit polyclonal to Rab 35.
Model	STJ95296
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, WB
Immunogen	Synthesized peptide derived from human Rab 35
Immunogen Region	70-150 aa, Internal
Gene ID	11021
Gene Symbol	RAB35
Dilution range	WB 1:500-1:2000ELISA 1:10000
Specificity	Rab 35 Polyclonal Antibody detects endogenous levels of Rab 35 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Ras-related protein Rab-35 GTP-binding protein RAY Ras-related protein Rab-1C
Molecular Weight	24 kDa
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:9774OMIM:604199
Alternative Names	Ras-related protein Rab-35 GTP-binding protein RAY Ras-related protein Rab-1C
Function	The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. That Rab is involved in the process of endocytosis and is an essential rate-limiting regulator of the fast recycling pathway back to the plasma membrane. During cytokinesis, required for the postfurling terminal steps, namely for intercellular bridge stability and abscission, possibly by controlling phosphatidylinositol 4,5-bis phosphate (PIP2) and SEPT2 localization at the intercellular bridge. May indirectly regulate neurite outgrowth. Together with TBC1D13 may be involved in regulation of insulin-induced glucose transporter SLC2A4/GLUT4 translocation to the plasma membrane in adipocytes.
Cellular Localization	Cell membrane Membrane, clathrin-coated pit Cytoplasmic vesicle, clathrin-coated vesicle Endosome Melanosome. Present on sorting endosomes and recycling endosome tubules . Tends to be enriched in PIP2-positive cell membrane domains . During mitosis, associated with the plasma membrane and present at the ingressing furrow during early cytokinesis as well as at the intercellular bridge later during cytokinesis . Identified in stage I to stage IV melanosomes .
Post-translational Modifications	AMPylation at Tyr-77 by L.pneumophila DrrA occurs in the switch 2 region and leads to moderate inactivation of the GTPase activity. It appears to prolong the lifetime of the GTP state of RAB1B by restricting access of GTPase effectors to switch 2 and blocking effector-stimulated GTP hydrolysis, thereby rendering RAB35 constitutively active. Phosphocholinated by L.pneumophila AnkX. Both GDP-bound and GTP-bound forms can be phosphocholinated. Phosphocholination inhibits the GEF activity of DENND1A.