


## Anti-Rad18 antibody



**Description** Rabbit polyclonal to Rad18.

Model STJ95324

**Host** Rabbit

**Reactivity** Human, Rat **Applications** ELISA, WB

**Immunogen** Synthesized peptide derived from human Rad18

**Immunogen Region** 270-350 aa, Internal

**Gene ID** <u>56852</u>

Gene Symbol RAD18

**Dilution range** WB 1:500-1:2000ELISA 1:40000

**Specificity** Rad18 Polyclonal Antibody detects endogenous levels of Rad18 protein.

**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Note** For Research Use Only (RUO).

Protein Name E3 ubiquitin-protein ligase RAD18 Postreplication repair protein RAD18

hHR18 hRAD18 RING finger protein 73 RING-type E3 ubiquitin transferase

RAD18

Molecular Weight 56 kDa

**Clonality** Polyclonal

**Conjugation** Unconjugated

**Isotype** IgG

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

**Concentration** 1 mg/ml

**Storage Instruction** Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:18278OMIM:605256</u>

Alternative Names E3 ubiquitin-protein ligase RAD18 Postreplication repair protein RAD18

hHR18 hRAD18 RING finger protein 73 RING-type E3 ubiquitin transferase

RAD18

**Function** E3 ubiquitin-protein ligase involved in postreplication repair of UV-damaged

DNA. Postreplication repair functions in gap-filling of a daughter strand on replication of damaged DNA. Associates to the E2 ubiquitin conjugating enzyme UBE2B to form the UBE2B-RAD18 ubiquitin ligase complex involved in mono-ubiquitination of DNA-associated PCNA on 'Lys-164'. Has

ssDNA binding activity.

Cellular Localization Nucleus Cytoplasm, cytoskeleton, microtubule organizing center, centrosome.

Associates with chromatin . Colocalizes with SLF1 in the nucleus and to centrosomes . Relocalizes with SLF1 to nuclear foci in response to DNA damage . Accumulates with the SLF1-SLF2 and SMC5-SMC6 complexes at replication-coupled DNA interstrand repair and DNA double-strand breaks

(DSBs) sites on chromatin in a ubiquitin-dependent manner .

St John's Laboratory Ltd

F +44 (0)207 681 2580

T+44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com