


Anti-Rad51C antibody





Description Rabbit polyclonal to Rad51C.

Model STJ95332

Host Rabbit

Reactivity Human, Simian

Applications ELISA, IF, IHC, WB

Immunogen Synthesized peptide derived from human Rad51C

Immunogen Region 130-210 aa, Internal

Gene ID <u>5889</u>

Gene Symbol RAD51C

Dilution range WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:5000

Specificity Rad51C Polyclonal Antibody detects endogenous levels of Rad51C protein.

Tissue Specificity Expressed in a variety of tissues, with highest expression in testis, heart

muscle, spleen and prostate.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name DNA repair protein RAD51 homolog 3 R51H3 RAD51 homolog C RAD51-

like protein 2

Molecular Weight 50 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. **Formulation**

1 mg/ml Concentration

Store at -20°C, and avoid repeat freeze-thaw cycles. **Storage Instruction**

Database Links HGNC:9820OMIM:602774

DNA repair protein RAD51 homolog 3 R51H3 RAD51 homolog C RAD51-**Alternative Names**

like protein 2

Function Essential for the homologous recombination (HR) pathway of DNA repair.

> Involved in the homologous recombination repair (HRR) pathway of doublestranded DNA breaks arising during DNA replication or induced by DNAdamaging agents. Part of the RAD21 paralog protein complexes BCDX2 and CX3 which act at different stages of the BRCA1-BRCA2-dependent HR pathway. Upon DNA damage, BCDX2 seems to act downstream of BRCA2 recruitment and upstream of RAD51 recruitment; CX3 seems to act downstream of RAD51 recruitment; both complexes bind predominantly to the intersection of the four duplex arms of the Holliday junction (HJ) and to junction of replication forks. The BCDX2 complex was originally reported to

bind single-stranded DNA, single-stranded gaps in duplex DNA and specifically to nicks in duplex DNA. The BCDX2 subcomplex

RAD51B:RAD51C exhibits single-stranded DNA-dependent ATPase activity suggesting an involvement in early stages of the HR pathway. Involved in

RAD51 foci formation in response to DNA damage suggesting an involvement in early stages of HR probably in the invasion step. Has an early function in DNA repair in facilitating phosphorylation of the checkpoint kinase CHEK2 and thereby transduction of the damage signal, leading to cell cycle arrest and HR activation. Participates in branch migration and HJ resolution and thus is important for processing HR intermediates late in the DNA repair process; the function may be linked to the CX3 complex. Part of a PALB2-scaffolded HR complex containing BRCA2 and which is thought to play a role in DNA repair by HR. Protects RAD51 from ubiquitin-mediated degradation that is enhanced following DNA damage. Plays a role in regulating mitochondrial DNA copy number under conditions of oxidative stress in the presence of RAD51 and XRCC3. Contributes to DNA cross-link resistance, sister chromatid cohesion and genomic stability. Involved in

maintaining centrosome number in mitosis.

Cellular Localization Nucleus. Cytoplasm. Cytoplasm, perinuclear region. Mitochondrion. DNA

> damage induces an increase in nuclear levels. Accumulates in DNA damage induced nuclear foci or RAD51C foci which is formed during the S or G2 phase of cell cycle. Accumulation at DNA lesions requires the presence of

NBN/NBS1, ATM and RPA.