

Anti-RELT antibody



Description	Rabbit polyclonal to RELT.
Model	STJ95411
Host	Rabbit
Reactivity	Human, Mouse
Applications	ELISA, IF, IHC, WB
Immunogen	Synthesized peptide derived from human RELT
Immunogen Region	350-430 aa, C-terminal
Gene ID	84957
Gene Symbol	RELT
Dilution range	WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:40000
Specificity	RELT Polyclonal Antibody detects endogenous levels of RELT protein.
Tissue Specificity	Highest levels are in spleen, lymph node, thymus, peripheral blood leukocytes, bone marrow and fetal liver. Very low levels in skeletal muscle, testis and colon. Not detected in brain, kidney and pancreas.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Tumor necrosis factor receptor superfamily member 19L Receptor expressed in lymphoid tissues
Molecular Weight	46 kDa

Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:13764OMIM:611211
Alternative Names	Tumor necrosis factor receptor superfamily member 19L Receptor expressed in lymphoid tissues
Function	Mediates activation of NF-kappa-B. May play a role in T-cell activation.
Cellular Localization	Cell membrane Cytoplasm
Post-translational Modifications	Phosphorylated in vitro by OXSR1.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com