


Anti-RFC3 antibody



Description	Rabbit polyclonal to RFC3.
Model	STJ95426
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human RFC3.
Immunogen Region	Internal
Gene ID	5983
Gene Symbol	RFC3
Dilution range	WB 1:500-1:2000IHC 1:100-1:300ELISA 1:40000
Specificity	RFC3 Polyclonal Antibody detects endogenous levels of RFC3 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Replication factor C subunit 3 Activator 1 38 kDa subunit A1 38 kDa subunit Activator 1 subunit 3 Replication factor C 38 kDa subunit RF-C 38 kDa subunit RFC38
Molecular Weight	40 kDa
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:9971OMIM:600405
Alternative Names	Replication factor C subunit 3 Activator 1 38 kDa subunit A1 38 kDa subunit Activator 1 subunit 3 Replication factor C 38 kDa subunit RF-C 38 kDa subunit RFC38
Function	The elongation of primed DNA templates by DNA polymerase delta and epsilon requires the action of the accessory proteins proliferating cell nuclear antigen (PCNA) and activator 1.
Cellular Localization	Nucleus

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com