


Anti-Ribosomal Protein L17 antibody



Description	Rabbit polyclonal to Ribosomal Protein L17.
Model	STJ95463
Host	Rabbit
Reactivity	Human, Mouse, Rat, Simian
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human Ribosomal Protein L17
Immunogen Region	70-150 aa, Internal
Gene ID	6139
Gene Symbol	RPL17
Dilution range	WB 1:500-1:2000IHC 1:100-1:300ELISA 1:10000
Specificity	Ribosomal Protein L17 Polyclonal Antibody detects endogenous levels of Ribosomal Protein L17 protein.
Tissue Specificity	Expressed in pancreas, lung, colon, cystic duct, gall bladder, kidney and liver. Expressed at high levels in the well differentiated pancreatic tumor cell lines HPAF, COLO 357 and Capan-1, the moderately differentiated pancreatic tumor cell lines T3M-4, AsPc-1 and BxPc-3, the poorly differentiated pancreatic tumor cell line MIA PaCa-2, and the pancreatic tumor cell lines of undefined differentiation status such as SW979. Expressed at lower levels in the poorly differentiated pancreatic tumor cell lines HCG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Note	For Research Use Only (RUO).
Protein Name	60S ribosomal protein L17 60S ribosomal protein L23 Large ribosomal subunit protein uL22 PD-1
Molecular Weight	24 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:10307OMIM:603661
Alternative Names	60S ribosomal protein L17 60S ribosomal protein L23 Large ribosomal subunit protein uL22 PD-1
Function	Component of the large ribosomal subunit.

St John's Laboratory Ltd

F +44 (0)207 681 2580
T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>
E info@stjohnslabs.com