


Anti-RNF130 antibody



Description	Rabbit polyclonal to RNF130.
Model	STJ95519
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, WB
Immunogen	Synthesized peptide derived from human RNF130
Immunogen Region	200-280 aa, C-terminal
Gene ID	55819
Gene Symbol	RNF130
Dilution range	WB 1:500-1:2000ELISA 1:5000
Specificity	RNF130 Polyclonal Antibody detects endogenous levels of RNF130 protein.
Tissue Specificity	Ubiquitously expressed. Highly expressed in leukocytes. Not expressed in erythroblasts.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	E3 ubiquitin-protein ligase RNF130 Goliath homolog H-Goliath RING finger protein 130 RING-type E3 ubiquitin transferase RNF130
Molecular Weight	36 kDa
Clonality	Polyclonal

Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:18280 OMIM:NA
Alternative Names	E3 ubiquitin-protein ligase RNF130 Goliath homolog H-Goliath RING finger protein 130 RING-type E3 ubiquitin transferase RNF130
Function	May have a role during the programmed cell death of hematopoietic cells . Acts as an E3 ubiquitin-protein ligase.
Cellular Localization	Membrane Cytoplasm
Post-translational Modifications	In vivo measurements suggest this protein is glycosylated . In contrast, in vitro experiments failed to detect glycosylation .

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com