

Anti-SRp20 antibody



Description Rabbit polyclonal to SRp20.

Model STJ95782

Host Rabbit

Reactivity Human, Mouse

Applications ELISA, IF, IHC, WB

Immunogen Synthesized peptide derived from human SRp20

Immunogen Region 80-160 aa, C-terminal

Gene ID <u>6428</u>

Gene Symbol SRSF3

Dilution range WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:5000

Specificity SRp20 Polyclonal Antibody detects endogenous levels of SRp20 protein.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name Serine/arginine-rich splicing factor 3 Pre-mRNA-splicing factor SRP20

Splicing factor, arginine/serine-rich 3

Molecular Weight 25 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Concentration 1 mg/ml

Storage Instruction Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links HGNC:107850MIM:603364

Alternative Names Serine/arginine-rich splicing factor 3 Pre-mRNA-splicing factor SRP20

Splicing factor, arginine/serine-rich 3

Function Splicing factor that specifically promotes exon-inclusion during alternative

splicing. Interaction with YTHDC1, a RNA-binding protein that recognizes

and binds N6-methyladenosine (m6A)-containing RNAs, promotes

recruitment of SRSF3 to its mRNA-binding elements adjacent to m6A sites, leading to exon-inclusion during alternative splicing . Also functions as export adapter involved in mRNA nuclear export such as of histone H2A . Binds mRNA which is thought to be transferred to the NXF1-NXT1 heterodimer for export (TAP/NXF1 pathway); enhances NXF1-NXT1 RNA-binding activity .

RNA-binding is semi-sequence specific.

Cellular Localization Nucleus Nucleus speckle Cytoplasm. Recruited to nuclear speckles following

interaction with YTHDC1.

Post-translational Phosphorylated by CLK1, CLK2, CLK3 and CLK4. Extensively

Modifications phosphorylated on serine residues in the RS domain.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com