


Anti-MMP-3 antibody



Description	Rabbit polyclonal to MMP-3.
--------------------	-----------------------------

Model	STJ95830
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, IF, IHC, WB
Immunogen	Synthesized peptide derived from human MMP-3
Immunogen Region	390-470 aa, C-terminal
Gene ID	4314
Gene Symbol	MMP3
Dilution range	WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:10000
Specificity	MMP-3 Polyclonal Antibody detects endogenous levels of MMP-3 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Stromelysin-1 SL-1 Matrix metalloproteinase-3 MMP-3 Transin-1
Molecular Weight	54 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG

Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:7173 OMIM:185250
Alternative Names	Stromelysin-1 SL-1 Matrix metalloproteinase-3 MMP-3 Transin-1
Function	Can degrade fibronectin, laminin, gelatins of type I, III, IV, and V; collagens III, IV, X, and IX, and cartilage proteoglycans. Activates procollagenase.
Sequence and Domain Family	The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.
Cellular Localization	Secreted, extracellular space, extracellular matrix

St John's Laboratory Ltd

F +44 (0)207 681 2580

W <http://www.stjohnslabs.com/>

T +44 (0)208 223 3081

E info@stjohnslabs.com