

Anti-TAF II p135/p105 antibody



Description	Rabbit polyclonal to TAF II p135/p105.
Model	STJ95892
Host	Rabbit
Reactivity	Human, Mouse
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human TAF II p135/p105
Immunogen Region	910-990 aa, C-terminal
Gene ID	6874
Gene Symbol	TAF4
Dilution range	WB 1:500-1:2000IHC 1:100-1:300ELISA 1:20000
Specificity	TAF II p135/p105 Polyclonal Antibody detects endogenous levels of TAF II p135/p105 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Transcription initiation factor TFIID subunit 4 RNA polymerase II TBP-associated factor subunit C TBP-associated factor 4 Transcription initiation factor TFIID 130 kDa subunit TAF II130 TAFII-130 TAFII130 Tr
Molecular Weight	110 kDa
Clonality	Polyclonal

Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:11537OMIM:601796
Alternative Names	Transcription initiation factor TFIID subunit 4 RNA polymerase II TBP-associated factor subunit C TBP-associated factor 4 Transcription initiation factor TFIID 130 kDa subunit TAF II130 TAFII-130 TAFII130 Tr
Function	Part of the TFIID complex, a multimeric protein complex that plays a central role in mediating promoter responses to various activators and repressors. Potentiates transcriptional activation by the AF-2S of the retinoic acid, vitamin D3 and thyroid hormone.
Cellular Localization	Nucleus.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com