

Anti-TAF II p18 antibody



Description	Rabbit polyclonal to TAF II p18.
Model	STJ95893
Host	Rabbit
Reactivity	Human, Mouse
Applications	ELISA, IHC
Immunogen	Synthesized peptide derived from human TAF II p18
Immunogen Region	40-120 aa, Internal
Gene ID	6884
Gene Symbol	TAF13
Dilution range	IHC 1:100-1:300ELISA 1:40000
Specificity	TAF II p18 Polyclonal Antibody detects endogenous levels of TAF II p18 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Transcription initiation factor TFIID subunit 13 Transcription initiation factor TFIID 18 kDa subunit TAF II18 TAFII-18 TAFII18
Molecular Weight	14.287 kDa
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:11546 OMIM:600774
Alternative Names	Transcription initiation factor TFIID subunit 13 Transcription initiation factor TFIID 18 kDa subunit TAF II18 TAFII-18 TAFII18
Function	TFIID beta-specific TAFII.
Sequence and Domain Family	The binding of TAF10 and TAF11 requires distinct domains of TAF13.
Cellular Localization	Nucleus.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com