

Anti-TAF II p250 antibody



Description Rabbit polyclonal to TAF II p250.

Model STJ95894

Host Rabbit

Reactivity Human, Mouse

Applications ELISA, IHC

Immunogen Synthesized peptide derived from human TAF II p250

Immunogen Region 1100-1180 aa, Internal

Gene ID <u>6872</u>

Gene Symbol TAF1

Dilution range IHC 1:100-1:300ELISA 1:10000

Specificity TAF II p250 Polyclonal Antibody detects endogenous levels of TAF II p250

protein.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name Transcription initiation factor TFIID subunit 1 Cell cycle gene 1 protein TBP-

associated factor 250 kDa p250 Transcription initiation factor TFIID 250 kDa

subunit TAF II250 TAFII-250 TAFII250

Molecular Weight 212.677 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Concentration 1 mg/ml

Storage Instruction Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:11535OMIM:300966</u>

Alternative Names Transcription initiation factor TFIID subunit 1 Cell cycle gene 1 protein TBP-

associated factor 250 kDa p250 Transcription initiation factor TFIID 250 kDa

subunit TAF II250 TAFII-250 TAFII250

Function Largest component and core scaffold of the TFIID basal transcription factor

complex . Contains novel N- and C-terminal Ser/Thr kinase domains which can autophosphorylate or transphosphorylate other transcription factors. Phosphorylates TP53 on 'Thr-55' which leads to MDM2-mediated degradation of TP53. Phosphorylates GTF2A1 and GTF2F1 on Ser residues. Possesses

DNA-binding activity . Essential for progression of the G1 phase of the cell cycle . Exhibits histone acetyltransferase activity towards histones H3 and H4

vice. Exhibits instone acceptualisterase activity towards instones 115 and 11-

Cellular Localization Nucleus

Post-translational

Modifications

Phosphorylated by casein kinase II in vitro.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com