


Anti-TBPL2 antibody



Description	Rabbit polyclonal to TBPL2.
Model	STJ95928
Host	Rabbit
Reactivity	Human
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human TBPL2
Immunogen Region	50-130 aa, Internal
Gene ID	387332
Gene Symbol	TBPL2
Dilution range	WB 1:500-1:2000IHC 1:100-1:300ELISA 1:5000
Specificity	TBPL2 Polyclonal Antibody detects endogenous levels of TBPL2 protein.
Tissue Specificity	Ubiquitously expressed in all tissues examined with highest levels in heart, lung, ovary, spleen and testes.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	TATA box-binding protein-like protein 2 TBP-like protein 2 TATA box-binding protein-related factor 3 TBP-related factor 3
Molecular Weight	41 kDa
Clonality	Polyclonal

Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:19841 OMIM:608964
Alternative Names	TATA box-binding protein-like protein 2 TBP-like protein 2 TATA box-binding protein-related factor 3 TBP-related factor 3
Function	Transcription factor required in complex with TAF3 for the differentiation of myoblasts into myocytes. The complex replaces TFIID at specific promoters at an early stage in the differentiation process .
Cellular Localization	Cytoplasm Nucleus. Present in the cytoplasm during cytokinesis.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com