


## Anti-TDAG51 antibody



<b>Description</b>	Rabbit polyclonal to TDAG51.
<b>Model</b>	STJ95955
<b>Host</b>	Rabbit
<b>Reactivity</b>	Human, Mouse, Rat
<b>Applications</b>	ELISA, WB
<b>Immunogen</b>	Synthesized peptide derived from human TDAG51
<b>Immunogen Region</b>	240-320 aa, C-terminal
<b>Gene ID</b>	<a href="#">22822</a>
<b>Gene Symbol</b>	<a href="#">PHLDA1</a>
<b>Dilution range</b>	WB 1:500-1:2000ELISA 1:10000
<b>Specificity</b>	TDAG51 Polyclonal Antibody detects endogenous levels of TDAG51 protein.
<b>Tissue Specificity</b>	Widely expressed with highest levels in pancreas. Strongly expressed by benign melanocytic nevi, and progressively reduced expressed in primary and metastatic melanomas (at protein level).
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Note</b>	For Research Use Only (RUO).
<b>Protein Name</b>	Pleckstrin homology-like domain family A member 1 Apoptosis-associated nuclear protein Proline- and glutamine-rich protein PQ-rich protein PQR protein Proline- and histidine-rich protein T-cell death-associated gene 51

<b>Molecular Weight</b>	40 kDa
<b>Clonality</b>	Polyclonal
<b>Conjugation</b>	Unconjugated
<b>Isotype</b>	IgG
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Concentration</b>	1 mg/ml
<b>Storage Instruction</b>	Store at -20°C, and avoid repeat freeze-thaw cycles.
<b>Database Links</b>	<a href="https://www.ncbi.nlm.nih.gov/Protein/89330">HGNC:89330</a> <a href="https://www.ncbi.nlm.nih.gov/Protein/MIM:605335">MIM:605335</a>
<b>Alternative Names</b>	Pleckstrin homology-like domain family A member 1 Apoptosis-associated nuclear protein Proline- and glutamine-rich protein PQ-rich protein PQR protein Proline- and histidine-rich protein T-cell death-associated gene 51
<b>Function</b>	Seems to be involved in regulation of apoptosis. May be involved in detachment-mediated programmed cell death. May mediate apoptosis during neuronal development. May be involved in regulation of anti-apoptotic effects of IGF1. May be involved in translational regulation.
<b>Cellular Localization</b>	Cytoplasm Cytoplasmic vesicle Nucleus, nucleolus. Colocalizes with intracellular vesicles.

---

**St John's Laboratory Ltd**

**F** +44 (0)207 681 2580

**T** +44 (0)208 223 3081

**W** <http://www.stjohnslabs.com/>

**E** [info@stjohnslabs.com](mailto:info@stjohnslabs.com)