

Anti-TEM7 antibody



Description Rabbit polyclonal to TEM7.

Model STJ95966

Host Rabbit

Reactivity Human

Applications ELISA, IHC

Immunogen Synthesized peptide derived from human TEM7

Immunogen Region 40-120 aa, Internal

Gene ID <u>57125</u>

Gene Symbol PLXDC1

Dilution range IHC 1:100-1:300ELISA 1:20000

Specificity TEM7 Polyclonal Antibody detects endogenous levels of TEM7 protein.

Tissue Specificity Detected in endothelial cells from colorectal cancer, and in endothelial cells

from primary cancers of the lung, liver, pancreas, breast and brain. Not detectable in endothelial cells from normal tissue. Expressed in fibrovascular membrane with increased expression in individuals with proliferative diabetic

retinopathy.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name Plexin domain-containing protein 1 Tumor endothelial marker 3 Tumor

endothelial marker 7

Molecular Weight 55.76 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Concentration 1 mg/ml

Storage Instruction Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links HGNC:209450MIM:606826

Alternative Names Plexin domain-containing protein 1 Tumor endothelial marker 3 Tumor

endothelial marker 7

Function Plays a critical role in endothelial cell capillary morphogenesis.

Cellular Localization Isoform 1: Cell membrane. Single-pass type I membrane protein. Cell

junction, tight junction. Localized predominantly at the tight junctions of vascular endothelial cells and to a lesser extent at the luminal surface of vascular endothelial cells.. Isoform 2: Secreted Isoform 3: Secreted Isoform 4:

Cytoplasm

Post-translational

Modifications

N-glycosylated.

St John's Laboratory Ltd

 \mathbf{F} +44 (0)207 681 2580

T +44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com