


Anti-TFDP1 antibody



Description Rabbit polyclonal to TFDP1.

Model STJ95976

Host Rabbit

Reactivity Human

Applications ELISA, IF, IHC, WB

Immunogen Synthesized peptide derived from human TFDP1

Immunogen Region 330-410 aa, C-terminal

Gene ID <u>7027</u>

Gene Symbol TFDP1

Dilution range WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:20000

Specificity TFDP1 Polyclonal Antibody detects endogenous levels of TFDP1 protein.

Tissue Specificity Highest levels in muscle. Also expressed in brain, placenta, liver and kidney.

Lower levels in lung and pancreas. Not detected in heart.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name Transcription factor Dp-1 DRTF1-polypeptide 1 DRTF1 E2F dimerization

partner 1

Molecular Weight 55 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. **Formulation**

1 mg/ml Concentration

Store at -20°C, and avoid repeat freeze-thaw cycles. **Storage Instruction**

Database Links HGNC:11749OMIM:189902

Transcription factor Dp-1 DRTF1-polypeptide 1 DRTF1 E2F dimerization **Alternative Names**

partner 1

Function Can stimulate E2F-dependent transcription. Binds DNA cooperatively with

> E2F family members through the E2 recognition site, 5'-TTTC[CG]CGC-3', found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication. The E2F1:DP complex appears to mediate both cell proliferation and apoptosis. Blocks adipocyte differentiation by repressing CEBPA binding to its target gene

promoters.

Cellular Localization Nucleus Cytoplasm. Shuttles between the cytoplasm and nucleus and

translocates into the nuclear compartment upon heterodimerization with E2F1.

Phosphorylation by E2F1-bound cyclin A-CDK2, in the S phase, inhibits E2F-Post-translational

mediated DNA binding and transactivation. Ubiquitinated by the **Modifications**

BCR(KBTBD5) complex, leading to its subsequent degradation.

St John's Laboratory Ltd

F +44 (0)207 681 2580

W http://www.stjohnslabs.com/ T+44 (0)208 223 3081 E info@stjohnslabs.com