


Anti-TFIIE-beta antibody



Description	Rabbit polyclonal to TFIIE-beta.
Model	STJ95981
Host	Rabbit
Reactivity	Human, Mouse
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human TFIIE-beta
Immunogen Region	120-200 aa, Internal
Gene ID	2961
Gene Symbol	GTF2E2
Dilution range	WB 1:500-1:2000IHC 1:100-1:300ELISA 1:20000
Specificity	TFIIE-beta Polyclonal Antibody detects endogenous levels of TFIIE-beta protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Transcription initiation factor IIE subunit beta TFIIE-beta General transcription factor IIE subunit 2
Molecular Weight	33 kDa
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:4651 OMIM:189964
Alternative Names	Transcription initiation factor IIE subunit beta TFIIE-beta General transcription factor IIE subunit 2
Function	Recruits TFIIH to the initiation complex and stimulates the RNA polymerase II C-terminal domain kinase and DNA-dependent ATPase activities of TFIIH. Both TFIIH and TFIIE are required for promoter clearance by RNA polymerase.
Cellular Localization	Nucleus

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com