

Anti-TFIIH p62 antibody



Description	Rabbit polyclonal to TFIIH p62.
Model	STJ95983
Host	Rabbit
Reactivity	Human, Mouse
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human TFIIH p62
Immunogen Region	10-90 aa, N-terminal
Gene ID	2965
Gene Symbol	GTF2H1
Dilution range	WB 1:500-1:2000IHC 1:100-1:300ELISA 1:20000
Specificity	TFIIH p62 Polyclonal Antibody detects endogenous levels of TFIIH p62 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	General transcription factor IIH subunit 1 Basic transcription factor 2 62 kDa subunit BTF2 p62 General transcription factor IIH polypeptide 1 TFIIH basal transcription factor complex p62 subunit
Molecular Weight	57 kDa
Clonality	Polyclonal

Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:4655OMIM:189972
Alternative Names	General transcription factor IIH subunit 1 Basic transcription factor 2 62 kDa subunit BTF2 p62 General transcription factor IIH polypeptide 1 TFIID basal transcription factor complex p62 subunit
Function	Component of the core-TFIID basal transcription factor involved in nucleotide excision repair (NER) of DNA and, when complexed to CAK, in RNA transcription by RNA polymerase II.
Cellular Localization	Nucleus.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com