


Anti-TFIIC110 antibody



Description	Rabbit polyclonal to TFIIC110.
Model	STJ95987
Host	Rabbit
Reactivity	Human
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human TFIIC110
Immunogen Region	40-120 aa, N-terminal
Gene ID	2976
Gene Symbol	GTF3C2
Dilution range	WB 1:500-1:2000IHC 1:100-1:300ELISA 1:20000
Specificity	TFIIC110 Polyclonal Antibody detects endogenous levels of TFIIC110 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	General transcription factor 3C polypeptide 2 TF3C-beta Transcription factor IIC 110 kDa subunit TFIIC 110 kDa subunit TFIIC110 Transcription factor IIC subunit beta
Molecular Weight	100 kDa
Clonality	Polyclonal

Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:4665OMIM:604883
Alternative Names	General transcription factor 3C polypeptide 2 TF3C-beta Transcription factor IIIC 110 kDa subunit TFIIC 110 kDa subunit TFIIC110 Transcription factor IIIC subunit beta
Function	Required for RNA polymerase III-mediated transcription. Component of TFIIC that initiates transcription complex assembly on tRNA and is required for transcription of 5S rRNA and other stable nuclear and cytoplasmic RNAs. May play a direct role in stabilizing interactions of TFIIC2 with TFIIC1.
Cellular Localization	Nucleus.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com