


Anti-TIEG2 antibody



Description	Rabbit polyclonal to TIEG2.
Model	STJ96018
Host	Rabbit
Reactivity	Human, Mouse
Applications	ELISA, IF, IHC, WB
Immunogen	Synthesized peptide derived from human TIEG2
Immunogen Region	1-80 aa, N-terminal
Gene ID	8462
Gene Symbol	KLF11
Dilution range	WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:20000
Specificity	TIEG2 Polyclonal Antibody detects endogenous levels of TIEG2 protein.
Tissue Specificity	Ubiquitous. Higher expression in erythroid cells.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Kruppel-like factor 11 Transforming growth factor-beta-inducible early growth response protein 2 TGFB-inducible early growth response protein 2 TIEG-2
Molecular Weight	55 kDa
Clonality	Polyclonal

Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:11811OMIM:603301
Alternative Names	Krueppel-like factor 11 Transforming growth factor-beta-inducible early growth response protein 2 TGFB-inducible early growth response protein 2 TIEG-2
Function	Transcription factor . Activates the epsilon- and gamma-globin gene promoters and, to a much lower degree, the beta-globin gene and represses promoters containing SP1-like binding inhibiting cell growth . Represses transcription of SMAD7 which enhances TGF-beta signaling . Induces apoptosis .
Cellular Localization	Nucleus

St John's Laboratory Ltd

F +44 (0)207 681 2580
T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>
E info@stjohnslabs.com