


Anti-TNF-R2 antibody



Description Rabbit polyclonal to TNF-R2.

Model STJ96053

Host Rabbit

Reactivity Human, Mouse, Rat

Applications ELISA, IF, IHC, WB

ImmunogenSynthesized peptide derived from human TNF-R2

Immunogen Region 350-430 aa, C-terminal

Gene ID <u>7133</u>

Gene Symbol TNFRSF1B

Dilution range WB 1:500-1:2000IHC 1:100-1:300IF 1:200-1:1000ELISA 1:20000

Specificity TNF-R2 Polyclonal Antibody detects endogenous levels of TNF-R2 protein.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name Tumor necrosis factor receptor superfamily member 1B Tumor necrosis factor

receptor 2 TNF-R2 Tumor necrosis factor receptor type II TNF-RII TNFR-II

p75 p80 TNF-alpha receptor CD antigen CD120b Etanerce

Molecular Weight 48 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Concentration 1 mg/ml

Storage Instruction Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links HGNC:11917OMIM:191191

Alternative Names Tumor necrosis factor receptor superfamily member 1B Tumor necrosis factor

receptor 2 TNF-R2 Tumor necrosis factor receptor type II TNF-RII TNFR-II

p75 p80 TNF-alpha receptor CD antigen CD120b Etanerce

Function Receptor with high affinity for TNFSF2/TNF-alpha and approximately 5-fold

lower affinity for homotrimeric TNFSF1/lymphotoxin-alpha. The TRAF1/TRAF2 complex recruits the apoptotic suppressors BIRC2 and BIRC3 to TNFRSF1B/TNFR2. This receptor mediates most of the metabolic effects of TNF-alpha. Isoform 2 blocks TNF-alpha-induced apoptosis, which suggests that it regulates TNF-alpha function by antagonizing its biological

activity.

Cellular Localization Isoform 1: Cell membrane. Single-pass type I membrane protein.. Isoform 2:

Secreted.. Tumor necrosis factor-binding protein 2: Secreted.

Post-translational Phosphorylated; mainly on serine residues and with a very low level on

Modifications threonine residues.; A soluble form (tumor necrosis factor binding protein 2)

is produced from the membrane form by proteolytic processing.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com