


Anti-TRADD antibody



Description	Rabbit polyclonal to TRADD.
Model	STJ96081
Host	Rabbit
Reactivity	Human, Mouse, Simian
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human TRADD
Immunogen Region	220-300 aa, C-terminal
Gene ID	8717
Gene Symbol	TRADD
Dilution range	WB 1:500-1:2000IHC 1:100-1:300ELISA 1:10000
Specificity	TRADD Polyclonal Antibody detects endogenous levels of TRADD protein.
Tissue Specificity	Found in all examined tissues.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Tumor necrosis factor receptor type 1-associated DEATH domain protein TNFR1-associated DEATH domain protein TNFRSF1A-associated via death domain
Molecular Weight	34 kDa
Clonality	Polyclonal

Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:12030OMIM:603500
Alternative Names	Tumor necrosis factor receptor type 1-associated DEATH domain protein TNFR1-associated DEATH domain protein TNFRSF1A-associated via death domain
Function	The nuclear form acts as a tumor suppressor by preventing ubiquitination and degradation of isoform p19ARF/ARF of CDKN2A by TRIP12: acts by interacting with TRIP12, leading to disrupt interaction between TRIP12 and isoform p19ARF/ARF of CDKN2A . Adapter molecule for TNFRSF1A/TNFR1 that specifically associates with the cytoplasmic domain of activated TNFRSF1A/TNFR1 mediating its interaction with FADD. Overexpression of TRADD leads to two major TNF-induced responses, apoptosis and activation of NF-kappa-B.
Sequence and Domain Family	Requires the intact death domain to associate with TNFRSF1A/TNFR1.
Cellular Localization	Nucleus Cytoplasm Cytoplasm, cytoskeleton. Shuttles between the cytoplasm and the nucleus.