


Anti-Translin antibody



Description	Rabbit polyclonal to Translin.
Model	STJ96087
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human Translin
Immunogen Region	70-150 aa, Internal
Gene ID	7247
Gene Symbol	TSN
Dilution range	WB 1:500-1:2000IHC 1:100-1:300ELISA 1:40000
Specificity	Translin Polyclonal Antibody detects endogenous levels of Translin protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Translin Component 3 of promoter of RISC C3PO
Molecular Weight	26 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG

Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:12379OMIM:600575
Alternative Names	Translin Component 3 of promoter of RISC C3PO
Function	DNA-binding protein that specifically recognizes consensus sequences at the breakpoint junctions in chromosomal translocations, mostly involving immunoglobulin (Ig)/T-cell receptor gene segments. Seems to recognize single-stranded DNA ends generated by staggered breaks occurring at recombination hot spots.; Exhibits both single-stranded and double-stranded endoribonuclease activity. May act as an activator of RNA-induced silencing complex (RISC) by facilitating endonucleolytic cleavage of the siRNA passenger strand.
Cellular Localization	Cytoplasm Nucleus

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com