


Anti-TReP-132 antibody



Description	Rabbit polyclonal to TReP-132.
Model	STJ96097
Host	Rabbit
Reactivity	Human, Mouse
Applications	ELISA, IHC
Immunogen	Synthesized peptide derived from human TReP-132
Immunogen Region	1040-1120 aa, C-terminal
Gene ID	55809
Gene Symbol	TRERF1
Dilution range	IHC 1:100-1:300ELISA 1:5000
Specificity	TReP-132 Polyclonal Antibody detects endogenous levels of TReP-132 protein.
Tissue Specificity	Highest expression was seen in thymus, testis and adrenal cortex, expressed also in the adrenal medulla, thyroid, and stomach. Highly expressed in steroidogenic JEG-3 and MCF-7 cells, low expression was seen in non-steroidogenic Hep-G2 and HEK293 cells.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Transcriptional-regulating factor 1 Breast cancer anti-estrogen resistance 2 Transcriptional-regulating protein 132 Zinc finger protein rapa Zinc finger

	transcription factor TReP-132
Molecular Weight	132.256 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:18273OMIM:610322
Alternative Names	Transcriptional-regulating factor 1 Breast cancer anti-estrogen resistance 2 Transcriptional-regulating protein 132 Zinc finger protein rapa Zinc finger transcription factor TReP-132
Function	Binds DNA and activates transcription of CYP11A1. Interaction with CREBBP and EP300 results in a synergistic transcriptional activation of CYP11A1.
Cellular Localization	Nucleus

St John's Laboratory Ltd

F +44 (0)207 681 2580
T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>
E info@stjohnslabs.com