

Anti-TSEN54 antibody



Description Rabbit polyclonal to TSEN54.

Model STJ96124

Host Rabbit

Reactivity Human

Applications ELISA, IHC, WB

Immunogen Synthesized peptide derived from human TSEN54

Immunogen Region 230-310 aa, Internal

Gene ID 283989

Gene Symbol TSEN54

Dilution range WB 1:500-1:2000IHC 1:100-1:300ELISA 1:20000

Specificity TSEN54 Polyclonal Antibody detects endogenous levels of TSEN54 protein.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name tRNA-splicing endonuclease subunit Sen54 SEN54 homolog HsSEN54

tRNA-intron endonuclease Sen54

Molecular Weight 60 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Concentration 1 mg/ml

Storage Instruction Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:27561OMIM:225753</u>

Alternative Names tRNA-splicing endonuclease subunit Sen54 SEN54 homolog HsSEN54

tRNA-intron endonuclease Sen54

Function Non-catalytic subunit of the tRNA-splicing endonuclease complex, a complex

responsible for identification and cleavage of the splice sites in pre-tRNA. It cleaves pre-tRNA at the 5' and 3' splice sites to release the intron. The products are an intron and two tRNA half-molecules bearing 2',3' cyclic phosphate and 5'-OH termini. There are no conserved sequences at the splice sites, but the intron is invariably located at the same site in the gene, placing the splice sites an invariant distance from the constant structural features of the tRNA body. The tRNA splicing endonuclease is also involved in mRNA processing via its association with pre-mRNA 3'-end processing factors, establishing a link between pre-tRNA splicing and pre-mRNA 3'-end formation, suggesting that the endonuclease subunits function in multiple

RNA-processing events.

Cellular Localization Nucleus Nucleus, nucleolus. May be transiently localized in the nucleolus.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com