


## Anti-USP50 antibody



|                         |                                                                                                                       |
|-------------------------|-----------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>      | Rabbit polyclonal to USP50.                                                                                           |
| <b>Model</b>            | STJ96209                                                                                                              |
| <b>Host</b>             | Rabbit                                                                                                                |
| <b>Reactivity</b>       | Human, Mouse                                                                                                          |
| <b>Applications</b>     | ELISA, IHC, WB                                                                                                        |
| <b>Immunogen</b>        | Synthesized peptide derived from human USP50                                                                          |
| <b>Immunogen Region</b> | 220-300 aa, C-terminal                                                                                                |
| <b>Gene ID</b>          | <a href="#">373509</a>                                                                                                |
| <b>Gene Symbol</b>      | <a href="#">USP50</a>                                                                                                 |
| <b>Dilution range</b>   | WB 1:500-1:2000IHC 1:100-1:300ELISA 1:20000                                                                           |
| <b>Specificity</b>      | USP50 Polyclonal Antibody detects endogenous levels of USP50 protein.                                                 |
| <b>Purification</b>     | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| <b>Note</b>             | For Research Use Only (RUO).                                                                                          |
| <b>Protein Name</b>     | Inactive ubiquitin carboxyl-terminal hydrolase 50 Inactive ubiquitin-specific peptidase 50                            |
| <b>Molecular Weight</b> | 38 kDa                                                                                                                |
| <b>Clonality</b>        | Polyclonal                                                                                                            |
| <b>Conjugation</b>      | Unconjugated                                                                                                          |

|                            |                                                                                            |
|----------------------------|--------------------------------------------------------------------------------------------|
| <b>Isotype</b>             | IgG                                                                                        |
| <b>Formulation</b>         | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.                    |
| <b>Concentration</b>       | 1 mg/ml                                                                                    |
| <b>Storage Instruction</b> | Store at -20°C, and avoid repeat freeze-thaw cycles.                                       |
| <b>Database Links</b>      | <a href="#">HGNC:20079</a> OMIM:NA                                                         |
| <b>Alternative Names</b>   | Inactive ubiquitin carboxyl-terminal hydrolase 50 Inactive ubiquitin-specific peptidase 50 |

---

**St John's Laboratory Ltd**

**F** +44 (0)207 681 2580

**T** +44 (0)208 223 3081

**W** <http://www.stjohnslabs.com/>

**E** [info@stjohnslabs.com](mailto:info@stjohnslabs.com)