


Anti-SAP 155 antibody



Description Rabbit polyclonal to SAP 155.

Model STJ96403

Host Rabbit

Reactivity Human, Mouse

Applications ELISA, WB

Immunogen Synthesized peptide derived from human SAP 155.

Immunogen Region N-terminal

Gene ID 23451

Gene Symbol SF3B1

Dilution range WB 1:500-1:2000ELISA 1:20000

Specificity SAP 155 Polyclonal Antibody detects endogenous levels of SAP 155 protein.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name Splicing factor 3B subunit 1 Pre-mRNA-splicing factor SF3b 155 kDa subunit

SF3b155 Spliceosome-associated protein 155 SAP 155

Molecular Weight 145 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Concentration 1 mg/ml

Store at -20°C, and avoid repeat freeze-thaw cycles. **Storage Instruction**

Database Links HGNC:10768OMIM:605590

Alternative Names Splicing factor 3B subunit 1 Pre-mRNA-splicing factor SF3b 155 kDa subunit

SF3b155 Spliceosome-associated protein 155 SAP 155

Function Subunit of the splicing factor SF3B required for 'A' complex assembly formed

> by the stable binding of U2 snRNP to the branchpoint sequence (BPS) in premRNA. Sequence independent binding of SF3A/SF3B complex upstream of the branch site is essential, it may anchor U2 snRNP to the pre-mRNA. May also be involved in the assembly of the 'E' complex. Belongs also to the minor U12-dependent spliceosome, which is involved in the splicing of rare class of

nuclear pre-mRNA intron.

Cellular Localization Nucleus speckle. During mitosis, transiently dispersed from the nuclear

speckles to the cytoplasm.

Post-translational Phosphorylated. Phosphorylation occurs concomitantly with the splicing **Modifications**

catalytic steps. Phosphorylation on Thr-244, Thr-248 and Thr-313 by cyclin-

dependent kinases promotes interaction with PPP1R8 during mitosis.

Citrullinated by PADI4.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T+44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com