


Anti-LIF antibody



Description	Rabbit polyclonal to LIF.
Model	STJ96472
Host	Rabbit
Reactivity	Human
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human LIF.
Immunogen Region	141-190 aa, C-terminal
Gene ID	3976
Gene Symbol	LIF
Dilution range	WB 1:500-1:2000IHC-P 1:100-300ELISA 1:20000
Specificity	LIF Polyclonal Antibody detects endogenous levels of LIF protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Leukemia inhibitory factor LIF Differentiation-stimulating factor D factor Melanoma-derived LPL inhibitor MLPLI Emfilermin
Molecular Weight	25 kDa
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:6596OMIM:159540
Alternative Names	Leukemia inhibitory factor LIF Differentiation-stimulating factor D factor Melanoma-derived LPL inhibitor MLPLI Emfilermin
Function	LIF has the capacity to induce terminal differentiation in leukemic cells. Its activities include the induction of hematopoietic differentiation in normal and myeloid leukemia cells, the induction of neuronal cell differentiation, and the stimulation of acute-phase protein synthesis in hepatocytes.
Cellular Localization	Secreted.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com