


Anti-Eotaxin-3 antibody



Description Rabbit polyclonal to Eotaxin-3.

Model STJ96515

Host Rabbit

Reactivity Human

Applications ELISA, IHC, WB

Immunogen Synthesized peptide derived from human Eotaxin-3.

Immunogen Region Internal

Gene ID <u>10344</u>

Gene Symbol <u>CCL26</u>

Dilution range WB 1:500-1:2000IHC-P 1:100-300ELISA 1:20000

Specificity Eotaxin-3 Polyclonal Antibody detects endogenous levels of Eotaxin-3

protein.

Tissue Specificity Ubiquitously expressed at low levels in various tissues including heart and

ovary.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name C-C motif chemokine 26 CC chemokine IMAC Eotaxin-3 Macrophage

inflammatory protein 4-alpha MIP-4-alpha Small-inducible cytokine A26

Thymic stroma chemokine-1 TSC-1

Molecular Weight 12 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Concentration 1 mg/ml

Storage Instruction Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:10625OMIM:604697</u>

Alternative Names C-C motif chemokine 26 CC chemokine IMAC Eotaxin-3 Macrophage

inflammatory protein 4-alpha MIP-4-alpha Small-inducible cytokine A26

Thymic stroma chemokine-1 TSC-1

Function Chemoattractant for eosinophils and basophils. Acts as a ligand for C-C

chemokine receptor CCR3 which triggers Ca(2+) mobilization in eosinophils .

Cellular Localization Secreted

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W http://www.stjohnslabs.com/

E info@stjohnslabs.com