


Anti-IL-16 antibody





Description	Rabbit polyclonal to IL-16.

Model STJ96540

Host Rabbit

Reactivity Human

Applications ELISA, IHC, WB

Immunogen Synthesized peptide derived from human IL-16.

Immunogen Region C-terminal

Gene ID 3603

Gene Symbol <u>IL16</u>

Dilution range WB 1:500-1:2000IHC-P 1:100-300ELISA 1:20000

Specificity IL-16 Polyclonal Antibody detects endogenous levels of IL-16 protein.

Tissue Specificity Isoform 3 is expressed in hemopoietic tissues, such as resting T-cells, but is

undetectable during active T-cell proliferation.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name Pro-interleukin-16 Interleukin-16 IL-16 Lymphocyte chemoattractant factor

LCF

Molecular Weight 66 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Concentration 1 mg/ml

Storage Instruction Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:5980OMIM:603035</u>

Alternative Names Pro-interleukin-16 Interleukin-16 IL-16 Lymphocyte chemoattractant factor

LCF

Function Interleukin-16 stimulates a migratory response in CD4+ lymphocytes,

monocytes, and eosinophils. Primes CD4+ T-cells for IL-2 and IL-15 responsiveness. Also induces T-lymphocyte expression of interleukin 2 receptor. Ligand for CD4.; Isoform 1 may act as a scaffolding protein that anchors ion channels in the membrane.; Isoform 3 is involved in cell cycle progression in T-cells. Appears to be involved in transcriptional regulation of SKP2 and is probably part of a transcriptional repression complex on the core promoter of the SKP2 gene. May act as a scaffold for GABPB1 (the DNA-binding subunit the GABP transcription factor complex) and HDAC3 thus maintaining transcriptional repression and blocking cell cycle progression in

resting T-cells.

Cellular Localization Interleukin-16: Secreted.. Isoform 1: Cytoplasm Isoform 3: Cytoplasm.

Nucleus.

Post-translational Isoform 3 is synthesized as a chemo-attractant inactive precursor in

Modifications hemopoietic tissues and is proteolytically cleaved by caspase-3 to yield IL-16.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com