


Anti-TF antibody



Description	Rabbit polyclonal to TF.
Model	STJ96559
Host	Rabbit
Reactivity	Human
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human TF.
Immunogen Region	131-180 aa, Internal
Gene ID	2152
Gene Symbol	F3
Dilution range	WB 1:500-1:2000IHC-P 1:100-300ELISA 1:20000
Specificity	TF Polyclonal Antibody detects endogenous levels of TF protein.
Tissue Specificity	Lung, placenta and pancreas.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Tissue factor TF Coagulation factor III Thromboplastin CD antigen CD142
Molecular Weight	33 kDa
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:3541OMIM:134390
Alternative Names	Tissue factor TF Coagulation factor III Thromboplastin CD antigen CD142
Function	Initiates blood coagulation by forming a complex with circulating factor VII or VIIa. The [TF:VIIa] complex activates factors IX or X by specific limited protolysis. TF plays a role in normal hemostasis by initiating the cell-surface assembly and propagation of the coagulation protease cascade.
Cellular Localization	Isoform 1: Membrane Isoform 2: Secreted

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com