


Anti-BLC antibody



Description	Rabbit polyclonal to BLC.
Model	STJ96573
Host	Rabbit
Reactivity	Human
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human BLC.
Immunogen Region	31-80 aa, Internal
Gene ID	10563
Gene Symbol	CXCL13
Dilution range	WB 1:500-1:2000IHC-P 1:100-300ELISA 1:20000
Specificity	BLC Polyclonal Antibody detects endogenous levels of BLC protein.
Tissue Specificity	Highest levels in liver, followed by spleen, lymph node, appendix and stomach. Low levels in salivary gland, mammary gland and fetal spleen.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	C-X-C motif chemokine 13 Angie B cell-attracting chemokine 1 BCA-1 B lymphocyte chemoattractant CXC chemokine BLC Small-inducible cytokine B13
Molecular Weight	12.664 kDa

Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:10639OMIM:605149
Alternative Names	C-X-C motif chemokine 13 Angiogenic cell-attracting chemokine 1 BCA-1 B lymphocyte chemoattractant CXC chemokine BLC Small-inducible cytokine B13
Function	Chemotactic for B-lymphocytes but not for T-lymphocytes, monocytes and neutrophils. Does not induce calcium release in B-lymphocytes. Binds to BLR1/CXCR5.
Cellular Localization	Secreted.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com