


## Anti-Protein C antibody



|                           |                                                                                                                                                                                                                     |
|---------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Description</b>        | Rabbit polyclonal to Protein C.                                                                                                                                                                                     |
| <b>Model</b>              | STJ96586                                                                                                                                                                                                            |
| <b>Host</b>               | Rabbit                                                                                                                                                                                                              |
| <b>Reactivity</b>         | Human                                                                                                                                                                                                               |
| <b>Applications</b>       | ELISA, IHC, WB                                                                                                                                                                                                      |
| <b>Immunogen</b>          | Synthesized peptide derived from human C protein                                                                                                                                                                    |
| <b>Immunogen Region</b>   | 181-230 aa, Internal                                                                                                                                                                                                |
| <b>Gene ID</b>            | <a href="#">5624</a>                                                                                                                                                                                                |
| <b>Gene Symbol</b>        | <a href="#">PROC</a>                                                                                                                                                                                                |
| <b>Dilution range</b>     | WB 1:500-1:2000IHC-P 1:100-300ELISA 1:20000                                                                                                                                                                         |
| <b>Specificity</b>        | Protein C Polyclonal Antibody detects endogenous levels of Protein C protein.                                                                                                                                       |
| <b>Tissue Specificity</b> | Plasma; synthesized in the liver.                                                                                                                                                                                   |
| <b>Purification</b>       | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.                                                                                               |
| <b>Note</b>               | For Research Use Only (RUO).                                                                                                                                                                                        |
| <b>Protein Name</b>       | Vitamin K-dependent protein C Anticoagulant protein C Autoprothrombin IIA<br>Blood coagulation factor XIV Vitamin K-dependent protein C light chain<br>Vitamin K-dependent protein C heavy chain Activation peptide |
| <b>Molecular Weight</b>   | 52 kDa                                                                                                                                                                                                              |
| <b>Clonality</b>          | Polyclonal                                                                                                                                                                                                          |

|                                         |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
|-----------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Conjugation</b>                      | Unconjugated                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| <b>Isotype</b>                          | IgG                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| <b>Formulation</b>                      | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| <b>Concentration</b>                    | 1 mg/ml                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| <b>Storage Instruction</b>              | Store at -20°C, and avoid repeat freeze-thaw cycles.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>Database Links</b>                   | <a href="#">HGNC:9451OMIM:176860</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>Alternative Names</b>                | Vitamin K-dependent protein C Anticoagulant protein C Autoprothrombin IIA<br>Blood coagulation factor XIV Vitamin K-dependent protein C light chain<br>Vitamin K-dependent protein C heavy chain Activation peptide                                                                                                                                                                                                                                                                                                                                                                                                       |
| <b>Function</b>                         | Protein C is a vitamin K-dependent serine protease that regulates blood coagulation by inactivating factors Va and VIIIa in the presence of calcium ions and phospholipids . Exerts a protective effect on the endothelial cell barrier function .                                                                                                                                                                                                                                                                                                                                                                        |
| <b>Cellular Localization</b>            | Secreted Golgi apparatus Endoplasmic reticulum                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| <b>Post-translational Modifications</b> | The vitamin K-dependent, enzymatic carboxylation of some Glu residues allows the modified protein to bind calcium.; N- and O-glycosylated. Partial (70%) N-glycosylation of Asn-371 with an atypical N-X-C site produces a higher molecular weight form referred to as alpha. The lower molecular weight form, not N-glycosylated at Asn-371, is beta. O-glycosylated with core 1 or possibly core 8 glycans. The iron and 2-oxoglutarate dependent 3-hydroxylation of aspartate and asparagine is (R) stereospecific within EGF domains. May be phosphorylated on a Ser or Thr in a region (AA 25-30) of the propeptide. |