


Anti-HDC antibody



Description	Rabbit polyclonal to HDC.
Model	STJ96590
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human HDC.
Immunogen Region	191-240 aa, Internal
Gene ID	3067
Gene Symbol	HDC
Dilution range	WB 1:500-1:2000IHC-P 1:100-300ELISA 1:20000
Specificity	HDC Polyclonal Antibody detects endogenous levels of HDC protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Histidine decarboxylase HDC
Molecular Weight	72/50 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG

Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:4855OMIM:142704
Alternative Names	Histidine decarboxylase HDC
Function	Catalyzes the biosynthesis of histamine from histidine.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com