


Anti-CD58 antibody



Description	Rabbit polyclonal to CD58.
Model	STJ96616
Host	Rabbit
Reactivity	Human
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human CD58.
Immunogen Region	21-70 aa, Internal
Gene ID	965
Gene Symbol	CD58
Dilution range	WB 1:500-1:2000IHC-P 1:100-300ELISA 1:20000
Specificity	CD58 Polyclonal Antibody detects endogenous levels of CD58 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Lymphocyte function-associated antigen 3 Ag3 Surface glycoprotein LFA-3 CD antigen CD58
Molecular Weight	27 kDa
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:1688OMIM:153420
Alternative Names	Lymphocyte function-associated antigen 3 Ag3 Surface glycoprotein LFA-3 CD antigen CD58
Function	Ligand of the T-lymphocyte CD2 glycoprotein. This interaction is important in mediating thymocyte interactions with thymic epithelial cells, antigen-independent and -dependent interactions of T-lymphocytes with target cells and antigen-presenting cells and the T-lymphocyte rosetting with erythrocytes. In addition, the LFA-3/CD2 interaction may prime response by both the CD2+ and LFA-3+ cells.
Cellular Localization	Isoform 1: Cell membrane. Single-pass type I membrane protein.. Isoform 2: Cell membrane. Lipid-anchor, GPI-anchor.