


Anti-Bcl-6 antibody



Western Blot (WB) analysis of 293 cells using Bcl-6 Antibody (STJ96676) from 2 batches.



Description

Bcl-6 is a protein encoded by the BCL6 gene which is approximately 78,8 kDa. Bcl-6 is localised to the nucleus. It is involved in cytokine signalling in immune system, gene expression and FoxO signalling pathway. The protein encoded by this gene is a zinc finger transcription factor and contains an N-terminal POZ domain. It is mainly required for germinal centre formation and antibody affinity maturation which has different mechanisms of action specific to the lineage and biological functions. It forms complexes with different corepressors and histone deacetylases to repress the transcriptional expression of different subsets of target genes. Bcl-6 is expressed in germinal centre T- and B-cells and in primary immature dendritic cells. Mutations in the BCL6 gene may result in b-cell lymphomas. STJ96676 was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. This polyclonal antibody detects endogenous levels of Bcl-6 protein.

Model	STJ96676
Host	Rabbit
Reactivity	Human, Mouse, Rat
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human Bcl-6.
Immunogen Region	271-320 aa, Internal
Gene ID	604
Gene Symbol	BCL6
Dilution range	WB 1:500-1:2000 IHC-P 1:100-1:300 ELISA 1:20000

Specificity	Bcl-6 Polyclonal Antibody detects endogenous levels of Bcl-6 protein.
Tissue Specificity	Expressed in germinal center T- and B-cells and in primary immature dendritic cells.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	B-cell lymphoma 6 protein BCL-6 B-cell lymphoma 5 protein BCL-5 Protein LAZ-3 Zinc finger and BTB domain-containing protein 27 Zinc finger protein 51
Molecular Weight	78 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:1001OMIM:109565
Alternative Names	B-cell lymphoma 6 protein BCL-6 B-cell lymphoma 5 protein BCL-5 Protein LAZ-3 Zinc finger and BTB domain-containing protein 27 Zinc finger protein 51
Function	Transcriptional repressor mainly required for germinal center (GC) formation and antibody affinity maturation which has different mechanisms of action specific to the lineage and biological functions. Forms complexes with different corepressors and histone deacetylases to repress the transcriptional expression of different subsets of target genes. Represses its target genes by binding directly to the DNA sequence 5'-TTCCTAGAA-3' (BCL6-binding site) or indirectly by repressing the transcriptional activity of transcription factors. In GC B-cells, represses genes that function in differentiation, inflammation, apoptosis and cell cycle control, also autoregulates its transcriptional expression and up-regulates, indirectly, the expression of some genes important for GC reactions, such as AICDA, through the repression of microRNAs expression, like miR155. An important function is to allow GC B-cells to proliferate very rapidly in response to T-cell dependent antigens and tolerate the physiological DNA breaks required for immunoglobulin class switch recombination and somatic hypermutation without inducing a p53/TP53-dependent apoptotic response. In follicular helper CD4(+) T-cells (T(FH) cells), promotes the expression of T(FH)-related genes but inhibits the differentiation of T(H)1, T(H)2 and T(H)17 cells. Also required for the establishment and maintenance of immunological memory for both T- and B-cells. Suppresses macrophage proliferation through competition with STAT5 for STAT-binding motifs binding on certain target genes, such as CCL2 and CCND2. In response to genotoxic stress, controls cell cycle arrest in GC B-cells in both p53/TP53-dependent and -independent manners. Besides, also controls neurogenesis through the alteration of the composition of NOTCH-dependent transcriptional complexes at selective NOTCH targets, such as

HES5, including the recruitment of the deacetylase SIRT1 and resulting in an epigenetic silencing leading to neuronal differentiation.

Sequence and Domain Family The BTB domain mediates homodimerization. Its dimer interface mediates peptide binding such as to corepressors BCOR and NCOR2 . Interaction with corepressors through the BTB domain is needed to facilitate the rapid proliferation and survival of GC B-cells but is not involved in the T(FH) formation and BCL6-mediated suppression of T(H)2 and T(H)17 differentiation required for GC formation .

Cellular Localization Nucleus

Post-translational Modifications Phosphorylated by MAPK1 in response to antigen receptor activation at Ser-333 and Ser-343. Phosphorylated by ATM in response to genotoxic stress. Phosphorylation induces its degradation by ubiquitin/proteasome pathway. Polyubiquitinated. Polyubiquitinated by SCF(FBXO11), leading to its degradation by the proteasome.; Acetylated at Lys-379 by EP300 which inhibits the interaction with NuRD complex and the transcriptional repressor function. Deacetylated by HDAC- and SIR2-dependent pathways.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com