
Anti-NKp44 antibody



Description Rabbit polyclonal to NKp44.

Model STJ96846

Host Rabbit

Reactivity Human

Applications ELISA, IHC, WB

Immunogen Synthesized peptide derived from human NKp44.

Immunogen Region 61-110 aa, Internal

Gene ID <u>9436</u>

Gene Symbol NCR2

Dilution range WB 1:500-1:2000IHC-P 1:100-1:300ELISA 1:20000

Specificity NKp44 Polyclonal Antibody detects endogenous levels of NKp44 protein.

Tissue Specificity Selectively expressed by activated NK cells and by in vitro cultured (i.e.

activated) TCRg/d lymphoid cells.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Note For Research Use Only (RUO).

Protein Name Natural cytotoxicity triggering receptor 2 Lymphocyte antigen 95 homolog

NK cell-activating receptor Natural killer cell p44-related protein NK-p44

NKp44 CD antigen CD336

Molecular Weight 44 kDa

Clonality Polyclonal

Conjugation Unconjugated

Isotype IgG

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Concentration 1 mg/ml

Storage Instruction Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links HGNC:6732OMIM:604531

Alternative Names Natural cytotoxicity triggering receptor 2 Lymphocyte antigen 95 homolog

NK cell-activating receptor Natural killer cell p44-related protein NK-p44

NKp44 CD antigen CD336

Function Cytotoxicity-activating receptor that may contribute to the increased

efficiency of activated natural killer (NK) cells to mediate tumor cell lysis.

Cellular Localization Cell membrane

St John's Laboratory Ltd

F +44 (0)207 681 2580

W http://www.stjohnslabs.com/ E info@stjohnslabs.com

T +44 (0)208 223 3081 **E** info@