


Anti-DcR1 antibody



Description	Rabbit polyclonal to DcR1.
Model	STJ96848
Host	Rabbit
Reactivity	Human
Applications	ELISA, WB
Immunogen	Synthesized peptide derived from human DcR1.
Immunogen Region	11-60 aa, Internal
Gene ID	8794
Gene Symbol	TNFRSF10C
Dilution range	WB 1:500-1:2000ELISA 1:20000
Specificity	DcR1 Polyclonal Antibody detects endogenous levels of DcR1 protein.
Tissue Specificity	Higher expression in normal tissues than in tumor cell lines. Highly expressed in peripheral blood lymphocytes, spleen, skeletal muscle, placenta, lung and heart.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	Tumor necrosis factor receptor superfamily member 10C Antagonist decoy receptor for TRAIL/Apo-2L Decoy TRAIL receptor without death domain Decoy receptor 1 DcR1 Lymphocyte inhibitor of TRAIL TNF-related apoptosis-induci

Molecular Weight	27 kDa
Clonality	Polyclonal
Conjugation	Unconjugated
Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:11906OMIM:603613
Alternative Names	Tumor necrosis factor receptor superfamily member 10C Antagonist decoy receptor for TRAIL/Apo-2L Decoy TRAIL receptor without death domain Decoy receptor 1 DcR1 Lymphocyte inhibitor of TRAIL TNF-related apoptosis-induci
Function	Receptor for the cytotoxic ligand TRAIL. Lacks a cytoplasmic death domain and hence is not capable of inducing apoptosis. May protect cells against TRAIL mediated apoptosis by competing with TRAIL-R1 and R2 for binding to the ligand.
Cellular Localization	Cell membrane. Lipid-anchor, GPI-anchor.
Post-translational Modifications	N-glycosylated and O-glycosylated.