


## Anti-AFP alpha 1 Fetoprotein antibody



Western Blot (WB) analysis of HepG2 cells using AFP alpha 1 Fetoprotein Monoclonal Antibody from two batches. (STJ96977)



**Description** 

AFP alpha 1 Fetoprotein is a protein encoded by the AFP gene which is approximately 68,6 kDa. AFP alpha 1 Fetoprotein is secreted into the extra cellular space. It is involved in hippo signalling pathway, embryonic stem cell differentiation pathways and FOXA2 and FOXA3 transcription factor networks. It is a major plasma protein produced by the yolk sac and the liver during fetal life. In adults, alpha-fetoprotein expression is often associated with hepatoma or teratoma and is thought to be the fetal counterpart of serum albumin. AFP alpha 1 Fetoprotein is expressed in the heart, liver, cells of the nervous system, lung and blood. Mutations in the AFP gene may result in an alpha-fetoprotein deficiency. STJ96977 was developed from clone 17C5 and was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. This antibody detects endogenous AFP proteins.

Model STJ96977

**Host** Mouse

**Reactivity** Human

**Applications** WB

Immunogen Synthetic Peptide

**Gene ID** <u>174</u>

Gene Symbol AFP

**Dilution range** WB 1:2000

**Specificity** The antibody detects endogenous AFP proteins.

**Tissue Specificity** Plasma. Synthesized by the fetal liver and yolk sac.

**Purification** The antibody was affinity-purified from mouse ascites by affinity-

chromatography using specific immunogen.

**Clone ID** 17C5

For Research Use Only (RUO). Note

Alpha-fetoprotein Alpha-1-fetoprotein Alpha-fetoglobulin **Protein Name** 

**Clonality** Monoclonal

Conjugation Unconjugated

IgG1 Isotype

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Store at -20°C, and avoid repeat freeze-thaw cycles. **Storage Instruction** 

**Database Links** HGNC:3170MIM:104150

**Alternative Names** Alpha-fetoprotein Alpha-1-fetoprotein Alpha-fetoglobulin

**Function** Binds copper, nickel, and fatty acids as well as, and bilirubin less well than,

serum albumin. Only a small percentage (less than 2%) of the human AFP

shows estrogen-binding properties.

**Cellular Localization** Secreted.

**Post-translational** Independent studies suggest heterogeneity of the N-terminal sequence of the

mature protein and of the cleavage site of the signal sequence.; Sulfated.

St John's Laboratory Ltd

**Modifications** 

**F** +44 (0)207 681 2580

W http://www.stjohnslabs.com/ T+44 (0)208 223 3081 E info@stjohnslabs.com