


## Anti-Ub (Acetyl K27) antibody



<b>Description</b>	Rabbit polyclonal to Acetyl-Ub (K27).
<b>Model</b>	STJ97356
<b>Host</b>	Rabbit
<b>Reactivity</b>	Human, Mouse, Rat
<b>Applications</b>	ELISA, IHC, WB
<b>Immunogen</b>	Synthesized peptide derived from the human Ub
<b>Immunogen Region</b>	around the acetylation site of K27.
<b>Gene ID</b>	<a href="#">7311</a>
<b>Gene Symbol</b>	<a href="#">UBA52</a>
<b>Dilution range</b>	WB 1:500-1:2000IHC-P 1:100-300ELISA 1:20000
<b>Specificity</b>	Acetyl-Ub (K27) Polyclonal AntibodySynthesized peptide derived from the human Ub around the acetylation site of K27.
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Note</b>	For Research Use Only (RUO).
<b>Protein Name</b>	Ubiquitin-60S ribosomal protein L40 CEP52 Ubiquitin A-52 residue ribosomal protein fusion product 1 Ubiquitin 60S ribosomal protein L40 Large ribosomal subunit protein eL40
<b>Clonality</b>	Polyclonal
<b>Conjugation</b>	Unconjugated

<b>Isotype</b>	IgG
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Concentration</b>	1 mg/ml
<b>Storage Instruction</b>	Store at -20°C, and avoid repeat freeze-thaw cycles.
<b>Database Links</b>	<a href="#">HGNC:12458</a> <a href="#">OMIM:191321</a>
<b>Alternative Names</b>	Ubiquitin-60S ribosomal protein L40 CEP52 Ubiquitin A-52 residue ribosomal protein fusion product 1 Ubiquitin 60S ribosomal protein L40 Large ribosomal subunit protein eL40
<b>Function</b>	Ubiquitin: Exists either covalently attached to another protein, or free (unanchored). When covalently bound, it is conjugated to target proteins via an isopeptide bond either as a monomer (monoubiquitin), a polymer linked via different Lys residues of the ubiquitin (polyubiquitin chains) or a linear polymer linked via the initiator Met of the ubiquitin (linear polyubiquitin chains). Polyubiquitin chains, when attached to a target protein, have different functions depending on the Lys residue of the ubiquitin that is linked: Lys-6-linked may be involved in DNA repair; Lys-11-linked is involved in ERAD (endoplasmic reticulum-associated degradation) and in cell-cycle regulation; Lys-29-linked is involved in lysosomal degradation; Lys-33-linked is involved in kinase modification; Lys-48-linked is involved in protein degradation via the proteasome; Lys-63-linked is involved in endocytosis, DNA-damage responses as well as in signaling processes leading to activation of the transcription factor NF-kappa-B. Linear polymer chains formed via attachment by the initiator Met lead to cell signaling. Ubiquitin is usually conjugated to Lys residues of target proteins, however, in rare cases, conjugation to Cys or Ser residues has been observed. When polyubiquitin is free (unanchored-polyubiquitin), it also has distinct roles, such as in activation of protein kinases, and in signaling.; 60S ribosomal protein L40: Component of the 60S subunit of the ribosome. Ribosomal protein L40 is essential for translation of a subset of cellular transcripts, and especially for cap-dependent translation of vesicular stomatitis virus mRNAs.
<b>Cellular Localization</b>	Ubiquitin: Cytoplasm Nucleus 60S ribosomal protein L40: Cytoplasm
<b>Post-translational Modifications</b>	Ubiquitin: Phosphorylated at Ser-65 by PINK1 during mitophagy. Phosphorylated ubiquitin specifically binds and activates parkin (PRKN), triggering mitophagy . Phosphorylation does not affect E1-mediated E2 charging of ubiquitin but affects discharging of E2 enzymes to form polyubiquitin chains. It also affects deubiquitination by deubiquitinase enzymes such as USP30 .