


## Anti-beta-tubulin antibody (HRP)



Western Blot (WB) analysis of HeLa cells using beta-tubulin (HRP Conjugated) Monoclonal Antibody diluted at 1:10,000; 40,000; 80,000; 320,000. (STJ97477)



### Description

Beta tubulin is a protein encoded by the tubb gene which is approximately 49,7 kDa. Beta tubulin is localised to the cytoskeleton and cytoplasm. It is involved in the regulation of PLK1 activity at G2/M transition, development of Slit-Robo signalling, the innate immune system, cell cycle and organelle biogenesis and maintenance. Beta tubulin contains a highly acidic C-terminal region which can bind cations such as calcium. Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha chain and forms part of the cytoskeleton. Beta tubulin is ubiquitously expressed in the spleen, thymus and immature brain. Mutations in the tubb gene result in complex cortical dysplasia with other brain malformations. Mutations can also cause congenital symmetric circumferential, an autosomal dominant disease which results in multiple rings of folded skin mostly affecting the limbs. STJ97477 was developed from clone 5G3. The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. The antibody detects endogenous beta tubulin (HRP Conjugated).

<b>Model</b>	STJ97477
<b>Host</b>	Mouse
<b>Reactivity</b>	Bovine, Canine, Hamster, Human, Insect, Mouse, Rabbit, Rat, Sheep, Simian, Yeast
<b>Applications</b>	IHC, WB
<b>Immunogen</b>	recombinant Protein
<b>Gene ID</b>	<a href="#">203068</a>
<b>Gene Symbol</b>	<a href="#">TUBB</a>

<b>Dilution range</b>	WB 1:1000-1:3000IHC 1:50-1:200
<b>Specificity</b>	The antibody detects beta-tubulin (HRP Conjugated)
<b>Tissue Specificity</b>	Ubiquitously expressed with highest levels in spleen, thymus and immature brain.
<b>Purification</b>	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
<b>Clone ID</b>	5G3
<b>Note</b>	For Research Use Only (RUO).
<b>Protein Name</b>	Tubulin beta chain Tubulin beta-5 chain
<b>Clonality</b>	Monoclonal
<b>Conjugation</b>	HRP
<b>Isotype</b>	IgG1
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Concentration</b>	1 mg/ml
<b>Storage Instruction</b>	Store at -20°C, and avoid repeat freeze-thaw cycles.
<b>Database Links</b>	<a href="#">HGNC:20778OMIM:156610</a>
<b>Alternative Names</b>	Tubulin beta chain Tubulin beta-5 chain
<b>Function</b>	Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha chain.
<b>Sequence and Domain Family</b>	The highly acidic C-terminal region may bind cations such as calcium.
<b>Cellular Localization</b>	Cytoplasm, cytoskeleton
<b>Post-translational Modifications</b>	Some glutamate residues at the C-terminus are polyglutamylated, resulting in polyglutamate chains on the gamma-carboxyl group . Polyglutamylation plays a key role in microtubule severing by spastin (SPAST). SPAST preferentially recognizes and acts on microtubules decorated with short polyglutamate tails: severing activity by SPAST increases as the number of glutamates per tubulin rises from one to eight, but decreases beyond this glutamylation threshold . Some glutamate residues at the C-terminus are monoglycylated but not polyglycylated due to the absence of functional TTLL10 in human. Monoglycylation is mainly limited to tubulin incorporated into axonemes (cilia and flagella). Both polyglutamylated and monoglycylated can coexist on the same protein on adjacent residues, and lowering glycylation levels increases polyglutamylated, and reciprocally. The precise function of monoglycylation is still unclear (Probable). Phosphorylated on Ser-172 by CDK1 during the cell cycle, from metaphase to telophase, but not in interphase. This phosphorylation inhibits tubulin incorporation into microtubules.