


Anti-MEK2 antibody



Description	Mouse monoclonal to MEK2 (2C3).
Model	STJ97518
Host	Mouse
Reactivity	Human, Mouse, Rat
Applications	WB
Immunogen	Synthesized peptide derived from human MEK-2 around the non-phosphorylation site of T394.
Gene ID	5605
Gene Symbol	MAP2K2
Dilution range	WB 1:1000-2000
Specificity	MEK2 Mouse Monoclonal Antibody (2C3) detects endogenous levels of MEK2
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Clone ID	2C3
Note	For Research Use Only (RUO).
Protein Name	Dual specificity mitogen-activated protein kinase kinase 2 MAP kinase kinase 2 MAPKK 2 ERK activator kinase 2 MAPK/ERK kinase 2 MEK 2
Clonality	Monoclonal
Conjugation	Unconjugated

Isotype	IgG1
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:68420MIM:601263
Alternative Names	Dual specificity mitogen-activated protein kinase kinase 2 MAP kinase kinase 2 MAPKK 2 ERK activator kinase 2 MAPK/ERK kinase 2 MEK 2
Function	Catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in MAP kinases. Activates the ERK1 and ERK2 MAP kinases .
Cellular Localization	Cytoplasm Membrane. Membrane localization is probably regulated by its interaction with KSR1.
Post-translational Modifications	MAPKK is itself dependent on Ser/Thr phosphorylation for activity catalyzed by MAP kinase kinase kinases (RAF or MEKK1). Phosphorylated by MAP2K1/MEK1 . Acetylation of Ser-222 and Ser-226 by Yersinia yopJ prevents phosphorylation and activation, thus blocking the MAPK signaling pathway.

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com