


Anti-CTACK antibody



Description

Rabbit polyclonal to CTACK.

Model	STJ97677
Host	Rabbit
Reactivity	Human
Applications	ELISA, IHC, WB
Immunogen	Synthesized peptide derived from human CTACK.
Immunogen Region	51-100 aa, Internal
Gene ID	10850
Gene Symbol	CCL27
Dilution range	WB 1:500-1:2000IHC-P 1:100-1:300ELISA 1:10000
Specificity	CTACK Polyclonal Antibody detects endogenous levels of CTACK protein.
Tissue Specificity	Testis, thymus, placenta, ovary and skin.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Note	For Research Use Only (RUO).
Protein Name	C-C motif chemokine 27 CC chemokine ILC Cutaneous T-cell-attracting chemokine CTACK ESkin IL-11 R-alpha-locus chemokine Skinkine Small-inducible cytokine A27
Clonality	Polyclonal
Conjugation	Unconjugated

Isotype	IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1 mg/ml
Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:10626 OMIM:604833
Alternative Names	C-C motif chemokine 27 CC chemokine ILC Cutaneous T-cell-attracting chemokine CTACK ESkin IL-11 R-alpha-locus chemokine Skinkine Small-inducible cytokine A27
Function	Chemotactic factor that attracts skin-associated memory T-lymphocytes. May play a role in mediating homing of lymphocytes to cutaneous sites. Binds to CCR10.
Cellular Localization	Secreted

St John's Laboratory Ltd

F +44 (0)207 681 2580

T +44 (0)208 223 3081

W <http://www.stjohnslabs.com/>

E info@stjohnslabs.com