


Anti-Beclin-1 antibody



| | |
|---------------------------|--|
| Description | Mouse monoclonal to Beclin-1. |
| Model | STJ97769 |
| Host | Mouse |
| Reactivity | Human, Mouse, Rat |
| Applications | IHC |
| Immunogen | synthetic peptide derived from Beclin-1 |
| Immunogen Region | 110-190 aa |
| Gene ID | 8678 |
| Gene Symbol | BECN1 |
| Dilution range | IHC 1:100-200 |
| Specificity | Beclin-1 Mouse Monoclonal Antibody (5A11) detects endogenous levels of BECN1 |
| Tissue Specificity | Ubiquitous. |
| Purification | The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. |
| Clone ID | 5A11 |
| Note | For Research Use Only (RUO). |
| Protein Name | Beclin-1 Coiled-coil myosin-like BCL2-interacting protein Protein GT197 Beclin-1-C 35 kDa Beclin-1-C 37 kDa |
| Clonality | Monoclonal |

| | |
|---|--|
| Conjugation | Unconjugated |
| Isotype | IgG1 |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Concentration | 1 mg/ml |
| Storage Instruction | Store at -20°C, and avoid repeat freeze-thaw cycles. |
| Database Links | HGNC:10340MIM:604378 |
| Alternative Names | Beclin-1 Coiled-coil myosin-like BCL2-interacting protein Protein GT197 Beclin-1-C 35 kDa Beclin-1-C 37 kDa |
| Function | Plays a central role in autophagy . Acts as core subunit of the PI3K complex that mediates formation of phosphatidylinositol 3-phosphate; different complex forms are believed to play a role in multiple membrane trafficking pathways: PI3KC3-C1 is involved in initiation of autophagosomes and PI3KC3-C2 in maturation of autophagosomes and endocytosis. Involved in regulation of degradative endocytic trafficking and required for the abscission step in cytokinesis, probably in the context of PI3KC3-C2 . Essential for the formation of PI3KC3-C2 but not PI3KC3-C1 PI3K complex forms. Involved in endocytosis . Protects against infection by a neurovirulent strain of Sindbis virus . May play a role in antiviral host defense. Beclin-1-C 35 kDa localized to mitochondria can promote apoptosis; it induces the mitochondrial translocation of BAX and the release of proapoptotic factors. |
| Sequence and Domain Family | The coiled coil domain can form antiparallel homodimers and mediates dimerization with the coiled coil domains of ATG14 or UVRAG involved in the formation of PI3K complexes. |
| Cellular Localization | Cytoplasm Golgi apparatus, trans-Golgi network membrane Endosome membrane Endoplasmic reticulum membrane Mitochondrion membrane Endosome Cytoplasmic vesicle, autophagosome. Interaction with ATG14 promotes translocation to autophagosomes. Expressed in dendrites and cell bodies of cerebellar Purkinje cells . Beclin-1-C 35 kDa: Mitochondrion Nucleus Cytoplasm Beclin-1-C 37 kDa: Mitochondrion |
| Post-translational Modifications | Phosphorylation at Thr-119 by DAPK1 reduces its interaction with BCL2 and BCL2L1 and promotes induction of autophagy . In response to autophagic stimuli, phosphorylated at serine residues by AMPK in an ATG14-dependent manner, and this phosphorylation is critical for maximally efficient autophagy . Polyubiquitinated by NEDD4, both with 'Lys11'- and 'Lys63'-linkages. 'Lys'-11-linked polyubiquitination leads to degradation and is enhanced when the stabilizing interaction partner VPS34 is depleted. Deubiquitinated by USP10 and USP13, leading to stabilize the PI3KC3/VPS34-containing complexes. Proteolytically processed by caspases including CASP8 and CASP3; the C-terminal fragments lack autophagy-inducing capacity and are proposed to induce apoptosis. Thus the cleavage is proposed to be a determinant to switch from autophagy to apoptosis pathways affecting cellular homeostasis including viral infections and survival of tumor cells. |