


## Anti-Smad3 antibody



<b>Description</b>	Mouse monoclonal to Smad3.
--------------------	----------------------------

<b>Model</b>	STJ97772
<b>Host</b>	Mouse
<b>Reactivity</b>	Human, Mouse, Rat
<b>Applications</b>	IHC, WB
<b>Immunogen</b>	synthetic peptide derived from Smad3
<b>Immunogen Region</b>	350-430 aa
<b>Gene ID</b>	<a href="#">4088</a>
<b>Gene Symbol</b>	<a href="#">SMAD3</a>
<b>Dilution range</b>	WB 1:1000-2000IHC 1:100-200
<b>Specificity</b>	Smad3 Mouse Monoclonal Antibody (4C9) detects endogenous levels of SMAD3
<b>Purification</b>	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
<b>Clone ID</b>	4C9
<b>Note</b>	For Research Use Only (RUO).
<b>Protein Name</b>	Mothers against decapentaplegic homolog 3 MAD homolog 3 Mad3 Mothers against DPP homolog 3 hMAD-3 JV15-2 SMAD family member 3 SMAD 3 Smad3 hSMAD3
<b>Clonality</b>	Monoclonal

<b>Conjugation</b>	Unconjugated
<b>Isotype</b>	IgG1
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Concentration</b>	1 mg/ml
<b>Storage Instruction</b>	Store at -20°C, and avoid repeat freeze-thaw cycles.
<b>Database Links</b>	<a href="#">HGNC:6769</a> <a href="#">OMIM:114500</a>
<b>Alternative Names</b>	Mothers against decapentaplegic homolog 3 MAD homolog 3 Mad3 Mothers against DPP homolog 3 hMAD-3 JV15-2 SMAD family member 3 SMAD 3 Smad3 hSMAD3
<b>Function</b>	Receptor-regulated SMAD (R-SMAD) that is an intracellular signal transducer and transcriptional modulator activated by TGF-beta (transforming growth factor) and activin type 1 receptor kinases. Binds the TRE element in the promoter region of many genes that are regulated by TGF-beta and, on formation of the SMAD3/SMAD4 complex, activates transcription. Also can form a SMAD3/SMAD4/JUN/FOS complex at the AP-1/SMAD site to regulate TGF-beta-mediated transcription. Has an inhibitory effect on wound healing probably by modulating both growth and migration of primary keratinocytes and by altering the TGF-mediated chemotaxis of monocytes. This effect on wound healing appears to be hormone-sensitive. Regulator of chondrogenesis and osteogenesis and inhibits early healing of bone fractures. Positively regulates PDPK1 kinase activity by stimulating its dissociation from the 14-3-3 protein YWHAQ which acts as a negative regulator.
<b>Sequence and Domain Family</b>	The MH1 domain is required for DNA binding. Also binds zinc ions which are necessary for the DNA binding.; The MH2 domain is required for both homomeric and heteromeric interactions and for transcriptional regulation. Sufficient for nuclear import.; The linker region is required for the TGFbeta-mediated transcriptional activity and acts synergistically with the MH2 domain.
<b>Cellular Localization</b>	Cytoplasm Nucleus. Cytoplasmic and nuclear in the absence of TGF-beta. On TGF-beta stimulation, migrates to the nucleus when complexed with SMAD4 . Through the action of the phosphatase PPM1A, released from the SMAD2/SMAD4 complex, and exported out of the nucleus by interaction with RANBP1 . Co-localizes with LEMD3 at the nucleus inner membrane . MAPK-mediated phosphorylation appears to have no effect on nuclear import . PDPK1 prevents its nuclear translocation in response to TGF-beta .
<b>Post-translational Modifications</b>	Phosphorylated on serine and threonine residues. Enhanced phosphorylation in the linker region on Thr-179, Ser-204 and Ser-208 on EGF and TGF-beta treatment. Ser-208 is the main site of MAPK-mediated phosphorylation. CDK-mediated phosphorylation occurs in a cell-cycle dependent manner and inhibits both the transcriptional activity and antiproliferative functions of SMAD3. This phosphorylation is inhibited by flavopiridol. Maximum phosphorylation at the G(1)/S junction. Also phosphorylated on serine residues in the C-terminal SXS motif by TGFBR1 and ACVR1. TGFBR1-mediated phosphorylation at these C-terminal sites is required for interaction with SMAD4, nuclear location and transactivational activity, and appears to be a prerequisite for the TGF-beta mediated phosphorylation in the linker region. Dephosphorylated in the C-terminal SXS motif by PPM1A. This

dephosphorylation disrupts the interaction with SMAD4, promotes nuclear export and terminates TGF-beta-mediated signaling. Phosphorylation at Ser-418 by CSNK1G2/CK1 promotes ligand-dependent ubiquitination and subsequent proteasome degradation, thus inhibiting SMAD3-mediated TGF-beta responses. Phosphorylated by PDPK1. Acetylation in the nucleus by EP300 in the MH2 domain regulates positively its transcriptional activity and is enhanced by TGF-beta. Poly-ADP-ribosylated by PARP1 and PARP2. ADP-ribosylation negatively regulates SMAD3 transcriptional responses during the course of TGF-beta signaling. Ubiquitinated. Monoubiquitinated, leading to prevent DNA-binding . Deubiquitination by USP15 alleviates inhibition and promotes activation of TGF-beta target genes . Ubiquitinated by RNF111, leading to its degradation: only SMAD3 proteins that are 'in use' are targeted by RNF111, RNF111 playing a key role in activating SMAD3 and regulating its turnover . Undergoes STUB1-mediated ubiquitination and degradation .

---

**St John's Laboratory Ltd****F** +44 (0)207 681 2580  
**T** +44 (0)208 223 3081**W** <http://www.stjohnslabs.com/>  
**E** [info@stjohnslabs.com](mailto:info@stjohnslabs.com)