


## Anti-c-Kit antibody



**Description** 

4

Mouse monoclonal to c-Kit.

Model STJ97958

**Host** Mouse

**Reactivity** Human

**Applications** ELISA, IHC, WB

**Immunogen** Purified recombinant fragment of c-Kit expressed in E. Coli.

**Gene ID** <u>3815</u>

Gene Symbol KIT

**Dilution range** WB 1:500-1:2000IHC 1:200-1:1000ELISA 1:10000

**Specificity** c-Kit Monoclonal Antibody detects endogenous levels of c-Kit protein.

**Tissue Specificity** Isoform 1 and isoform 2 are detected in spermatogonia and Leydig cells.

Isoform 3 is detected in round spermatids, elongating spermatids and spermatozoa (at protein level). Widely expressed. Detected in the

hematopoietic system, the gastrointestinal system, in melanocytes and in germ

cells.

**Purification** Affinity purification

Clone ID 8D7

**Note** For Research Use Only (RUO).

Protein Name Mast/stem cell growth factor receptor Kit SCFR Piebald trait protein PBT

Proto-oncogene c-Kit Tyrosine-protein kinase Kit p145 c-kit v-kit Hardy-

Zuckerman 4 feline sarcoma viral oncogene homolog CD antigen C

**Clonality** Monoclonal

**Conjugation** Unconjugated

Isotype IgG1

**Modifications** 

**Formulation** Ascitic fluid containing 0.03% sodium azide.

**Storage Instruction** Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:6342OMIM:164920</u>

Alternative Names Mast/stem cell growth factor receptor Kit SCFR Piebald trait protein PBT

Proto-oncogene c-Kit Tyrosine-protein kinase Kit p145 c-kit v-kit Hardy-

Zuckerman 4 feline sarcoma viral oncogene homolog CD antigen C

**Function** Tyrosine-protein kinase that acts as cell-surface receptor for the cytokine

KITLG/SCF and plays an essential role in the regulation of cell survival and proliferation, hematopoiesis, stem cell maintenance, gametogenesis, mast cell development, migration and function, and in melanogenesis. In response to

KITLG/SCF binding, KIT can activate several signaling pathways.

Phosphorylates PIK3R1, PLCG1, SH2B2/APS and CBL. Activates the AKT1 signaling pathway by phosphorylation of PIK3R1, the regulatory subunit of phosphatidylinositol 3-kinase. Activated KIT also transmits signals via GRB2 and activation of RAS, RAF1 and the MAP kinases MAPK1/ERK2 and/or MAPK3/ERK1. Promotes activation of STAT family members STAT1,

STAT3, STAT5A and STAT5B. Activation of PLCG1 leads to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate. KIT signaling is modulated by protein phosphatases, and by

rapid internalization and degradation of the receptor. Activated KIT promotes phosphorylation of the protein phosphatases PTPN6/SHP-1 and PTPRU, and of the transcription factors STAT1, STAT3, STAT5A and STAT5B. Promotes

phosphorylation of PIK3R1, CBL, CRK (isoform Crk-II), LYN, MAPK1/ERK2 and/or MAPK3/ERK1, PLCG1, SRC and SHC1.

**Cellular Localization** Isoform 1: Cell membrane. Single-pass type I membrane protein.. Isoform 2:

Cell membrane. Single-pass type I membrane protein.. Isoform 3: Cytoplasm. Detected in the cytoplasm of spermatozoa, especially in the equatorial and

subacrosomal region of the sperm head.

Post-translational Ubiquitinated by SOCS6. KIT is rapidly ubiquitinated after

autophosphorylation induced by KITLG/SCF binding, leading to

internalization and degradation. Autophosphorylated on tyrosine residues. KITLG/SCF binding enhances autophosphorylation. Isoform 1 shows low levels of tyrosine phosphorylation in the absence of added KITLG/SCF (in vitro). Kinase activity is down-regulated by phosphorylation on serine residues by protein kinase C family members. Phosphorylation at Tyr-568 is required for interaction with PTPN11/SHP-2, CRK (isoform Crk-II) and members of the SRC tyrosine-protein kinase family. Phosphorylation at Tyr-570 is required for interaction with PTPN6/SHP-1. Phosphorylation at Tyr-703, Tyr-823 and Tyr-936 is important for interaction with PIK3R1. Phosphorylation at Tyr-823 and Tyr-936 is important for interaction with

GRB7.

St John's Laboratory Ltd

**F** +44 (0)207 681 2580 **T** +44 (0)208 223 3081

W http://www.stjohnslabs.com/
E info@stjohnslabs.com