


Anti-c-Rel antibody



Description	Mouse monoclonal to c-Rel.
Model	STJ97966
Host	Mouse
Reactivity	Human, Mouse
Applications	ELISA, IF, IHC, WB
Immunogen	Purified recombinant fragment of human c-Rel expressed in E. Coli.
Gene ID	5966
Gene Symbol	REL
Dilution range	WB 1:500-1:2000IHC 1:200-1:1000IF 1:200-1:1000ELISA 1:10000
Specificity	c-Rel Monoclonal Antibody detects endogenous levels of c-Rel protein.
Purification	Affinity purification
Clone ID	10000000
Note	For Research Use Only (RUO).
Protein Name	Proto-oncogene c-Rel
Clonality	Monoclonal
Conjugation	Unconjugated
Isotype	IgG1
Formulation	Ascitic fluid containing 0.03% sodium azide.

Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:9954OMIM:164910
Alternative Names	Proto-oncogene c-Rel
Function	Proto-oncogene that may play a role in differentiation and lymphopoiesis. NF-kappa-B is a pleiotropic transcription factor which is present in almost all cell types and is involved in many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFkB1/p105, NFkB1/p50, REL and NFkB2/p52. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct preferences for different kappa-B sites that they can bind with distinguishable affinity and specificity. Different dimer combinations act as transcriptional activators or repressors, respectively. NF-kappa-B is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NF-kappa-B complexes are held in the cytoplasm in an inactive state complexed with members of the NF-kappa-B inhibitor (I-kappa-B) family. In a conventional activation pathway, I-kappa-B is phosphorylated by I-kappa-B kinases (IKKs) in response to different activators, subsequently degraded thus liberating the active NF-kappa-B complex which translocates to the nucleus. The NF-kappa-B heterodimer RELA/p65-c-Rel is a transcriptional activator.
Cellular Localization	Nucleus