


## Anti-EphB4 antibody



Description

Mouse monoclonal to EphB4.

Model STJ98036

**Host** Mouse

**Reactivity** Human

**Applications** ELISA, IF, WB

Immunogen Purified recombinant extracellular fragment of human EphB4 fused with

hIgGFc tag expressed in HEK293 cell line.

**Gene ID** <u>2050</u>

Gene Symbol <u>EPHB4</u>

**Dilution range** WB 1:500-1:2000IF 1:200-1:1000ELISA 1:10000

**Specificity** EphB4 Monoclonal Antibody detects endogenous levels of EphB4 protein.

**Tissue Specificity** Abundantly expressed in placenta but also detected in kidney, liver, lung,

pancreas, skeletal muscle and heart. Expressed in primitive and myeloid, but not lymphoid, hematopoietic cells. Also observed in cell lines derived from

liver, breast, colon, lung, melanocyte and cervix.

**Purification** Affinity purification

Clone ID 5B5

**Note** For Research Use Only (RUO).

**Protein Name** Ephrin type-B receptor 4 Hepatoma transmembrane kinase Tyrosine-protein

kinase TYRO11

**Clonality** Monoclonal

**Conjugation** Unconjugated

**Isotype** IgG1

**Formulation** Ascitic fluid containing 0.03% sodium azide.

**Storage Instruction** Store at -20°C, and avoid repeat freeze-thaw cycles.

Database Links <u>HGNC:3395OMIM:600011</u>

Alternative Names Ephrin type-B receptor 4 Hepatoma transmembrane kinase Tyrosine-protein

kinase TYRO11

**Function** Receptor tyrosine kinase which binds promiscuously transmembrane ephrin-B

family ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Together with its cognate ligand/functional ligand EFNB2 plays a central role in heart morphogenesis and angiogenesis through regulation of cell adhesion and cell migration. EPHB4-mediated forward signaling controls cellular repulsion and segregation form EFNB2-expressing cells. Plays also a role in postnatal blood vessel remodeling, morphogenesis and permeability

and is thus important in the context of tumor angiogenesis.

Cellular Localization Cell membrane

Post-translational

**Modifications** 

Phosphorylated; autophosphorylation is stimulated by EFNB2.

St John's Laboratory Ltd

**F** +44 (0)207 681 2580

T +44 (0)208 223 3081

W http://www.stjohnslabs.com/ E info@stjohnslabs.com