


Anti-GTBP antibody



Description	Mouse monoclonal to GTBP.
Model	STJ98124
Host	Mouse
Reactivity	Human
Applications	ELISA, WB
Immunogen	Purified recombinant fragment of human GTBP expressed in E. Coli.
Gene ID	2956
Gene Symbol	MSH6
Dilution range	WB 1:500-1:2000ELISA 1:10000
Specificity	GTBP Monoclonal Antibody detects endogenous levels of GTBP protein.
Purification	Affinity purification
Clone ID	5B11
Note	For Research Use Only (RUO).
Protein Name	DNA mismatch repair protein Msh6 hMSH6 G/T mismatch-binding protein GTBP GTMBP MutS protein homolog 6 MutS-alpha 160 kDa subunit p160
Clonality	Monoclonal
Conjugation	Unconjugated
Isotype	IgG1
Formulation	Ascitic fluid containing 0.03% sodium azide.

Storage Instruction	Store at -20°C, and avoid repeat freeze-thaw cycles.
Database Links	HGNC:7329OMIM:114500
Alternative Names	DNA mismatch repair protein Msh6 hMSH6 G/T mismatch-binding protein GTBP GTMBP MutS protein homolog 6 MutS-alpha 160 kDa subunit p160
Function	Component of the post-replicative DNA mismatch repair system (MMR). Heterodimerizes with MSH2 to form MutS alpha, which binds to DNA mismatches thereby initiating DNA repair. When bound, MutS alpha bends the DNA helix and shields approximately 20 base pairs, and recognizes single base mismatches and dinucleotide insertion-deletion loops (IDL) in the DNA. After mismatch binding, forms a ternary complex with the MutL alpha heterodimer, which is thought to be responsible for directing the downstream MMR events, including strand discrimination, excision, and resynthesis. ATP binding and hydrolysis play a pivotal role in mismatch repair functions. The ATPase activity associated with MutS alpha regulates binding similar to a molecular switch: mismatched DNA provokes ADP-->ATP exchange, resulting in a discernible conformational transition that converts MutS alpha into a sliding clamp capable of hydrolysis-independent diffusion along the DNA backbone. This transition is crucial for mismatch repair. MutS alpha may also play a role in DNA homologous recombination repair. Recruited on chromatin in G1 and early S phase via its PWWP domain that specifically binds trimethylated 'Lys-36' of histone H3 (H3K36me3): early recruitment to chromatin to be replicated allowing a quick identification of mismatch repair to initiate the DNA mismatch repair reaction.
Sequence and Domain Family	The PWWP domain specifically recognizes and binds trimethylated 'Lys-36' of histone H3 (H3K36me3).
Cellular Localization	Nucleus Chromosome. Associates with H3K36me3 via its PWWP domain.
Post-translational Modifications	The N-terminus is blocked.; Phosphorylated by PRKCZ, which may prevent MutS alpha degradation by the ubiquitin-proteasome pathway.